

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
24 April 2003 (24.04.2003)

PCT

(10) International Publication Number
WO 03/033514 A1

- (51) International Patent Classification⁷: **C07H 21/04**, A61K 48/00
- (21) International Application Number: PCT/IL02/00339
- (22) International Filing Date: 1 May 2002 (01.05.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/330,118 19 October 2001 (19.10.2001) US
- (71) Applicant (for all designated States except US): **VASCULAR BIOGENICS LTD.** [IL/IL]; 3 Habossem Street, 77610 Ashdod (IL).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **HARATS, Dror** [IL/IL]; 71 Mendes Street, Kiryat Krinitzy, 53765 Ramat Gan (IL). **GREENBERGER, Shoshana** [IL/IL]; 15 Hainbal Street, 49773 Petah Tikva (IL).
- (74) Agent: **G. E. EHRLICH (1995) LTD.**; 28 Bezalel Street, 52521 Ramat Gan (IL).
- (81) Designated States (*national*): AE, AG, AL, AM, AT (utility model), AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ (utility model), CZ, DE (utility model), DE, DK (utility model), DK, DM, DZ, EC, EE (utility model), EE, ES, FI (utility model), FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK (utility model), SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— with international search report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

WO 03/033514 A1

(54) Title: POLYNUCLEOTIDE CONSTRUCTS, PHARMACEUTICAL COMPOSITIONS AND METHODS FOR TARGETED DOWNREGULATION OF ANGIOGENESIS AND ANTICANCER THERAPY

(57) Abstract: A novel nucleic acid construct for down-regulating angiogenesis in a tissue of a subject is provided. The nucleic acid construct includes: (a) a first polynucleotide region encoding a chimeric polypeptide including a ligand binding domain fused to an effector domain of an apoptosis signaling molecule; and (b) a second polynucleotide region encoding a cis acting regulatory element being for directing expression of the chimeric polypeptide in a specific tissue or cell; wherein the ligand binding domain is selected such that it is capable of binding a ligand present in the specific tissue or cell, whereas binding of the ligand to the ligand binding domain activates the effector domain of the apoptosis signaling molecule. Also provided are methods of utilizing this nucleic acid construct for treating diseases characterized by excessive or aberrant neo-vascularization or cell growth.

POLYNUCLEOTIDE CONSTRUCTS, PHARMACEUTICAL
COMPOSITIONS AND METHODS FOR TARGETED
DOWNREGULATION OF ANGIOGENESIS AND ANTICANCER
THERAPY

5

FIELD AND BACKGROUND OF THE INVENTION

The present invention relates to a nucleic acid constructs,
pharmaceutical compositions and methods which can be used to
downregulate angiogenesis in specific tissue regions of a subject. More
10 particularly, the present invention relates to nucleic acid constructs, which can
be used to activate apoptosis in specific cell subsets, thus, enabling treatment
of diseases characterized by excessive or aberrant neovascularization or cell
growth.

Angiogenesis is the growth of new blood vessels, a process that
15 depends mainly on locomotion, proliferation, and tube formation by capillary
endothelial cells. During angiogenesis, endothelial cells emerge from their
quiescent state and proliferate rapidly. Although the molecular mechanisms
responsible for transition of a cell to angiogenic phenotype are not known, the
sequence of events leading to the formation of new vessels has been well
20 documented [Hanahan, D., Science 277, 48-50, (1997)]. The vascular growth
entails either endothelial sprouting [Risau, W., Nature 386, 671-674, (1997)]
or intussusceptions [Patan, S., et al; Microvasc. Res. 51, 260-272, (1996)]. In
the first pathway, the following sequence of events may occur: (a) dissolution
of the basement of the vessel, usually a post capillary venule, and the
25 interstitial matrix; (b) migration of endothelial cells toward the stimulus; (c)
proliferation of endothelial cells trailing behind the leading endothelial cell
(s); (d) formation of lumen (canalization) in the endothelial array/sprout; (e)
formation of branches and loops by confluential anastomoses of sprouts to
permit blood flow; (f) investment of the vessel with pericytes (i.e.,
30 periendothelial cells and smooth muscle cells); and (g) formation of basement
membrane around the immature vessel. New vessels can also be formed via

the second pathway: insertion of interstitial tissue columns into the lumen of preexisting vessels. The subsequent growth of these columns and their stabilization result in partitioning of the vessel lumen and remodeling of the local vascular network.

5 A variety of angiogenic factors govern the angiogenic process. It is understood that during pathology, the fine balance between pro-angiogenic factors and anti-angiogenic factors is disrupted, thereby eliciting nonself-limiting endothelial and periendothelial cell-proliferation. Until recently, the angiogenesis that occurs in diseases of ocular neovascularization, arthritis,
10 skin diseases, and tumors, had been difficult to suppress therapeutically.

Therefore, the fundamental goal of all anti-angiogenic therapy is to return foci of proliferating microvessels to their normal resting state, and to prevent their regrowth [Cancer: Principles & Practice of Oncology, Fifth Edition, edited by Vincent T. DeVita, Jr., Samuel Hellman, Steven A.
15 Rosenberg. Lippincott-Raven Publishers, Philadelphia. (1997)].

Anti-angiogenic therapy is a robust clinical approach, as it can delay the progression of tumor growth (e.g., retinopathies, benign and malignant angiogenic tumors).

In general, every disease caused by uncontrolled growth of capillary
20 blood vessels such as diabetic retinopathy, psoriasis, arthritis, hemangiomas, tumor growth and metastasis is a target for anti-angiogenic therapy.

For example, the progressive growth of solid tumors beyond clinically occult sizes (e.g., a few mm³) requires the continuous formation of new blood vessels, a process known as tumor angiogenesis. Tumor growth and
25 metastasis are angiogenesis-dependent. A tumor must continuously stimulate the growth of new capillary blood vessels to deliver nutrients and oxygen for the tumor itself to grow. Therefore, either prevention of tumor angiogenesis or selective destruction of tumor's existing blood vessels (vascular targeting therapy) underlies anti-angiogenic tumor therapy.

Recently, a plethora of anti-angiogenic agents has been developed for the treatment of malignant diseases, some of which are already under clinical trials (for review see Herbst et al. (2002) *Semin. Oncol.* 29:66-77).

The most studied target for tumor anti-angiogenic treatment is the dominant process regulating angiogenesis in human i.e., the interaction of vascular endothelial growth factor (VEGF) with its receptor (VEGFR). Agents which regulate VEGFR pro-angiogenic action include (i) antibodies directed at the VEGF protein itself or to the receptor (e.g., rhuMAb VEGF); (ii) small molecule compounds directed to the VEGFR tyrosine kinase (e.g., ZD6474 and SU5416); (iii) VEGFR targeted ribozymes.

Other novel angiogenesis inhibitors include 2-Methoxyestradiol (2-ME2) a natural metabolite of estradiol that possesses unique anti-tumor and anti-angiogenic properties and angiostatin and endostatin - proteolytic cleavage fragments of plasminogen and collagen XVIII, respectively.

Though promising in pre-clinical models, to date systemic administration of all anti-angiogenic agents tested in clinical trials, have shown limited rate of success and considerable toxicities including thrombocytopenia, leukopenia and hemoptysis. These results suggest that there may be limits to the use of current tumor anti-angiogenic agents as therapy for advanced malignancies. O'Reilly et al. have shown that the latency between the initiation of anti-angiogenic therapy and antitumor effect may result in initial tumor progression before response to therapy [O'Reilly S et al. (1998) *Proc Am Soc Clin Oncol* 17:217a]. Furthermore, recent studies suggest that the regulation of angiogenesis may differ among capillary beds, suggesting that anti-angiogenic therapy may need to be optimized on an organ/tissue-specific basis [Arap et al. (1998) *Science* 279:377-380].

Interestingly, poor results have also been obtained when anti-angiogenic therapy (e.g., heparin, heparin-peptide treatment) directed at smooth muscle cell proliferation has been practiced on myocardial ischemia in patients with coronary artery disease [Liu et al., *Circulation*, 79: 1374-1387

(1989); Goldman et al., *Atherosclerosis*, 65: 215-225 (1987); Wolinsky et al., *JACC*, 15 (2): 475-481 (1990)]. Various limitations associated with the use of such agents for the treatment of cardiovascular diseases included: (i) systemic toxicity creating intolerable level of risk for patients with cardiovascular diseases; (ii) interference with vascular wound healing following surgery; (iii) possible damage to surrounding endothelium and/or other medial smooth muscle cells.

In-light of these and the inherent obstacles associated with systemic administration of anti-angiogenic factors (i.e., manufacturing limitations based on in-vitro instability and high doses required; and peak kinetics of bolus administration attributing to sub-optimal effects) limit the effective use of angiogenic factors in treating neo-vascularization associated diseases.

With the identification of new genes that regulate the angiogenic process, somatic gene therapy has been attempted to overcome these limitations. Although, great efforts have been directed towards developing methods for gene therapy of cancer, cardiovascular and peripheral vascular diseases, there is still major obstacles to effective and specific gene delivery [for review see, Feldman AL. (2000) *Cancer* 89(6):1181-94] In general, the main limiting factor of gene therapy with a gene of interest, using a recombinant viral vector as a shuttle is the ability to specifically direct the gene of interest to the target tissue.

Attempts to overcome these limitations included the use of tissue-specific promoters conjugated to cytotoxic genes. For example, endothelial cell targeting of a cytotoxic gene, expressed under the control endothelial-specific promoters has been described by Jagger et al who used the KDR or E-selectin promoter to express TNF α specifically in endothelial cells [Jagger RT. Et al. *Hum Gene Ther* (1997) 8(18):2239-47]. Ozaki et al used the von-Willebrand factor (vWF) promoter to deliver herpes simplex virus thymidine kinase (HSV-tk) to HUVEC [*Hum Gene Ther* (1996) 7(13):1483-90].

However, these promoters showed only weak activity and did not allow for high levels of expression.

An alternate approach presented by Kong and Crystal included a tumor specific expression of anti-angiogenic factors. To date, however, the toxicity of recombinant forms of endogenous anti-angiogenic agents has not been demonstrated although some synthetic anti-angiogenic agents have been associated with toxicity in preclinical models [Kong and Crystal (1998) J. Natl. Cancer Inst. 90:273-76].

Angiostatin has also been used as a possible anti-angiogenic agent (Falkman et al, Cell 1997 Jan 24;88(2):277-85), however due to the redundancy of factors involved in regulation of angiogenesis in tumors, it is highly unlikely that angiostatin therapy alone would be effective.

There is thus a widely recognized need for, and it would be highly advantageous to have a novel approach for efficiently down-regulating angiogenesis in specific tissue regions of a subject while being devoid of the toxic side effects characterizing prior art anti-angiogenesis approaches.

SUMMARY OF THE INVENTION

According to one aspect of the present invention there is provided a nucleic acid construct comprising: (a) a first polynucleotide region encoding a chimeric polypeptide including a ligand binding domain fused to an effector domain of an apoptosis signaling molecule; and (b) a second polynucleotide region encoding a cis acting regulatory element being for directing expression of the chimeric polypeptide in a specific tissue or cell; wherein the ligand binding domain is selected such that it is capable of binding a ligand present in the specific tissue or cell, whereas binding of the ligand to the ligand binding domain activates the effector domain of the apoptosis signaling molecule.

According to further features in preferred embodiments of the invention described below, there is provided a mammalian cell transformed with the nucleic acid construct described above.

According to another aspect of the present invention there is provided
5 a method of down-regulating angiogenesis in a tissue of a subject, the method comprising administering to the subject a nucleic acid construct designed and configured for generating apoptosis in a sub-population of angiogenic cells, the nucleic acid construct including: (a) a first polynucleotide region encoding a chimeric polypeptide including a ligand binding domain fused to an effector
10 domain of an apoptosis signaling molecule; and (b) a second polynucleotide region encoding a cis acting regulatory element being for directing expression of the chimeric polypeptide in the sub-population of angiogenic cells; wherein the ligand binding domain is selected such that it is capable of binding a ligand present in, or provided to, the sub-population of angiogenic cells,
15 whereas binding of the ligand to the ligand binding domain activates the effector domain of the apoptosis signaling molecule, thereby down-regulating angiogenesis in the tissue.

According to further features in preferred embodiments of the invention described below, the method further comprising administering the
20 ligand to the subject in a manner suitable for providing the ligand to the sub-population of angiogenic cells.

According to yet another aspect of the present invention there is provided method of down-regulating angiogenesis in a tissue of a subject, the method comprising: (a) administering to the subject a nucleic acid construct
25 designed and configured for generating apoptosis in a sub-population of angiogenic cells, the nucleic acid construct including: (i) a first polynucleotide region encoding a chimeric polypeptide including a ligand binding domain fused to an effector domain of an apoptosis signaling molecule, wherein the effector domain is selected such that it is activated
30 following binding of a ligand to the ligand binding domain; and (ii) a second

polynucleotide region encoding a cis acting regulatory element being for directing expression of the chimeric polypeptide in the sub-population of angiogenic cells; and (b) administering to the subject the ligand, thereby down-regulating angiogenesis in the tissue.

5 According to still further features in the described preferred embodiments the administering the ligand to the subject is effected by a method selected from the group consisting of: (i) systemic in-vivo administration; (ii) ex-vivo administration to cells removed from a body of the subject, the cells subsequently reintroduced into the body of the subject;
10 and (iii) local in-vivo administration.

According to still further features in the described preferred embodiments the cis-acting regulatory element is an endothelial cell-specific or periendothelial cell-specific promoter selected from the group consisting of the PPE-1 promoter, the PPE-1-3x promoter, the TIE-1 promoter, the TIE-2
15 promoter, the Endoglin promoter, the von Willerband promoter, the KDR/flk-1 promoter, The FLT-1 promoter, the Egr-1 promoter, the ICAM-1 promoter, the VCAM-1 promoter, the PECAM-1 promoter, the CArG box element and aortic carboxypeptidase-like protein (ACLP) promoter.

According to still further features in the described preferred
20 embodiments the ligand binding domain is a ligand-binding domain of a cell-surface receptor.

According to still further features in the described preferred embodiments the cell-surface receptor is selected from the group consisting of a receptor tyrosine kinase, a receptor serine kinase, a receptor threonine
25 kinase, a cell adhesion molecule and a phosphatase receptor.

According to still further features in the described preferred embodiments the apoptosis signaling molecule is selected from the group consisting of Fas and TNFR.

According to still another aspect of the present invention there is
30 provided a pharmaceutical composition for down regulating angiogenesis in a

tissue of a subject, the pharmaceutical composition comprising as an active ingredient a nucleic acid construct designed and configured for generating apoptosis in a subpopulation of angiogenic cells and a pharmaceutical acceptable carrier, the nucleic acid construct including: (a) a first polynucleotide region encoding a chimeric polypeptide including a ligand binding domain fused to an effector domain of an apoptosis signaling molecule; and (b) a second polynucleotide region encoding a cis acting regulatory element being for directing expression of the chimeric polypeptide in the subpopulation of angiogenic cells; wherein the ligand binding domain is selected such that it is capable of binding a ligand present in the specific tissue or cell, whereas binding of the ligand to the ligand binding domain activates the effector domain of the apoptosis signaling molecule.

According to an additional aspect of the present invention there is provided a method of treating a disease or condition associated with excessive neo-vascularization, the method comprising administering a therapeutically effective amount of a nucleic acid construct designed and configured for generating apoptosis in a sub-population of angiogenic cells, the nucleic acid construct including: (i) a first polynucleotide region encoding a chimeric polypeptide including a ligand binding domain fused to an effector domain of an apoptosis signaling molecule; and (ii) a second polynucleotide region encoding a cis acting regulatory element being for directing expression of the chimeric polypeptide in the sub-population of angiogenic cells; wherein the ligand binding domain is selected such that it is capable of binding a ligand present in, or provided to, the sub-population of angiogenic cells, whereas binding of the ligand to the ligand binding domain activates the effector domain of the apoptosis signaling molecule, thereby down-regulating angiogenesis in the tissue and treating the disease or condition associated with excessive neo-vascularization.

According to yet an additional aspect of the present invention there is provided a method of treating a tumor in a subject, the method comprising

administering a therapeutically effective amount of a nucleic acid construct designed and configured for generating apoptosis in cells of the tumor, the nucleic acid construct including: (i) a first polynucleotide region encoding a chimeric polypeptide including a ligand binding domain fused to an effector domain of an apoptosis signaling molecule; and (ii) a second polynucleotide region encoding a cis acting regulatory element being for directing expression of the chimeric polypeptide in the cells of the tumor; wherein the ligand binding domain is selected such that it is capable of binding a ligand present in, or provided to, the cells of the tumor, whereas binding of the ligand to the ligand binding domain activates the effector domain of the apoptosis signaling molecule to thereby direct apoptosis in the cells of the tumor.

According to still further features in the described preferred embodiments the method further comprising administering the ligand to the subject in a manner suitable for providing the ligand to the cells of the tumor.

According to still further features in the described preferred embodiments the cis acting regulatory element is selected from the group consisting of the gastrin-releasing peptide (GRP) promoter, the hTERT promoter, the Hexokinase type II promoter and the L-plastin promoter.

The present invention successfully addresses the shortcomings of the presently known configurations by providing a novel approach for efficiently downregulating angiogenesis and tumor cell proliferation in a specific and targeted manner.

BRIEF DESCRIPTION OF THE DRAWINGS

The invention is herein described, by way of example only, with reference to the accompanying drawings. With specific reference now to the drawings in detail, it is stressed that the particulars shown are by way of example and for purposes of illustrative discussion of the preferred embodiments of the present invention only, and are presented in the cause of

providing what is believed to be the most useful and readily understood description of the principles and conceptual aspects of the invention. In this regard, no attempt is made to show structural details of the invention in more detail than is necessary for a fundamental understanding of the invention, the
5 description taken with the drawings making apparent to those skilled in the art how the several forms of the invention may be embodied in practice.

In the drawings:

FIGs. 1a-b are schematic illustrations of Fas chimera gene constructed from the extracellular region of TNFR1 and the trans-membrane and
10 intracellular regions of Fas and cloned into pcDNA3 plasmid (a) or into adenoviral vectors (b).

FIGs. 2a-b illustrate apoptotic activity of the pro-apoptotic genes, Fas chimera and TNFR1. Figure 2a - illustrates Bovine Aortic Endothelial Cells (BAEC) transfected with either pcDNA-3-TNFR1 (lower panel) or control
15 empty vector (upper panel) and an expression plasmid encoding GFP. Figure 2b - illustrates 293 Cells transfected with either pcDNA-3-Fas-c (lower panel) or control empty vector (upper panel) and an expression plasmid encoding GFP. Transfected cells were visualized using fluorescence microscopy and apoptotic activity was morphologically determined.

20 FIGs. 3a-f are electron microscopy images of BAEC cells transfected with pro-apoptotic genes. 24 hours post transfection, BAEC cells were fixed in 2.5% glutaraldehyde and processed. Presented are cells in successive stages of the apoptotic process.

FIG. 4 are histograms quantifying apoptotic activity of the indicated
25 pro-apoptotic genes in transfected BAEC and 293 cells.

FIG. 5a represents a PCR analysis of AdPPE-Fas-c. Lanes 1-2 - PCR products obtained using primers encompassing the PPE-1 promoter and Fas-c gene. Lanes 3-4 - PCR products obtained using Fas-c primers. Lanes 5-6 - PCR products obtained in the absence of template DNA.

FIG. 5b is a western blot analysis of AdPPE-Fas-c transfected BAEC cells. Protein samples were resolved by SDS-PAGE, transferred to nitrocellulose membrane and probed with a polyclonal antibody directed against the extracellular portion of TNFR1. Lane 1-2 - pcDNA3-Fas-c BAEC
5 transfected cells (positive control). Lane 3-4 - BAEC cells transfected with the indicated MOI of AdPPE-Fas-c viruses. Lane 5 - non-transfected cells. Lane 6-7 - BAEC cells transfected with the indicated MOI of AdPPE-Luc.

FIGs. 6a-d are photomicrographs illustrating the effect of Fas-chimera over-expression on apoptosis of endothelial cells. BAEC cells were infected
10 with: Ad-PPE-1-3x-Fas-chimera (Figure 6a); Ad-PPE-1-3x-luciferase (Figure 6b); Ad-PPE-1-3x-Fas-chimera and Ad-PPE1-3x-GFP (Figure 6c); Ad-PPE-1-3x-luciferase and Ad-PPE-1-3x-GFP; each at MOI 1000 (Figure 6d). Photomicrographs were taken 72 h post infection at x10 magnification.

FIG. 7 is a histogram illustrating apoptotic specific effect of Ad-PPE-
15 1-3x-Fas-chimera on endothelial cells. Viability of endothelial (BAEC, HUVEC) and non-endothelial (Normal skin fibroblasts-NSF) cells was quantified by crystal violet staining 72 h post infection with either Ad-PPE-1-3x-Fas-chimera or control (luciferase) virus.

FIG. 8 shows a dose response effect of TNF α administration on Fas-chimera mediated apoptosis. BAEC were infected with Ad-PPE-1-3x-Fas-c.
20 48 h post infection TNF was added to the growth medium (at the indicated dose). Viability was determined by the crystal violet assay 24 h thereafter.

FIGs. 9a-e are photomicrographs illustrating an endothelial cell-specific apoptosis mediated by the cooperative action of TNF α ligand and
25 Fas-c receptor. The indicated cells were incubated in the presence or absence of TNF α (10 ng/ml) 48 h following infection with Ad-PPE-1-3x-Fas-c; crystal violet staining was effected 72 h post infection.

FIG. 10a is a dose response curve illustrating the TNF α -dependent apoptotic effect of Ad-CMV-Fas-c on endothelial cells. Viability of BAEC

cells infected with the indicated MOI of Ad-CMV-Fas-chimera was determined following incubation with TNF α .

FIGs. 10b-d illustrate the apoptotic effect of TNF α ligand and Ad-CMV-Fas-chimera on the non-endothelial cells NSF. Figure 10b - NSF infected with a control virus. Figure 10c - NSF infected with Ad-CMV-Fas-chimera. Figure 10d - NSF infected with Ad-CMV-Fas-chimera and incubated with TNF (10 ng/ml).

FIGs. 11 a-c illustrate the *In-vivo* anti-tumoral effect of Ad-PPE-1-3x-Fas-c. Mice inoculated with B16 melanoma cells were injected intravenously with Ad-PPE-1-3x-Fas-c, Ad-CMV-Fas-chimera, control virus or saline when tumor was palpable. Figure 11a - tumor areas, measured during treatment period. Fig 11b - tumor weights at end of treatment period. Fig 11c - an image representing the state of the tumor in the Ad-PPE-1-3x-Fas-c treated mouse and the control mouse.

15

DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention is of nucleic acid constructs and methods, which can be used to treat diseases characterized by excessive or aberrant neovascularization or cell growth. Specifically, the present invention can be used to specifically target and kill cells involved in angiogenesis, thus enabling down-regulation of angiogenesis and anti-tumor therapy.

The principles and operation of the present invention may be better understood with reference to the drawings and accompanying descriptions.

Before explaining at least one embodiment of the invention in detail, it is to be understood that the invention is not limited in its application to the details of construction and the arrangement of the components set forth in the following description or illustrated in the drawings. The invention is capable of other embodiments or of being practiced or carried out in various ways. Also, it is to be understood that the phraseology and terminology employed

25

herein is for the purpose of description and should not be regarded as limiting.

One of the most fundamental goals of anti-angiogenic therapy is to return foci of proliferating microvessels to their normal resting state, and to prevent their re-growth.

To date, anti-angiogenic therapy approaches, which employ systemic administration of anti-angiogenic agents have been employed with limited success mostly due to the toxic side effects which lead to the formation of thrombocytopenia, leukopenia and/or hemoptysis in the treated individual. The toxic side effects associated with prior art approaches is a result of non-specific expression of the anti-angiogenic agents employed and exposure of healthy tissue to these agents or the redundancy and thus non-effectiveness of the anti-angiogenic agents used.

While reducing the present invention to practice, the present inventors have uncovered that a combination of tissue-specific expression and specific activation of a pro-apoptotic agent enables selective apoptosis of cells involved in angiogenesis without exposing non-targeted tissue or cells to these agents, thus, avoiding the toxic side effects and redundancy characterizing prior art treatment approaches.

Thus, according to one aspect of the present invention there is provided a method of down-regulating angiogenesis in a tissue of a subject. As used herein, the phrase "down-regulating angiogenesis" refers to either slowing down or stopping the angiogenic process, which lead to formation of new blood vessels.

The method according to this aspect of the present invention is effected by administering to the subject a nucleic acid construct designed and configured for generating apoptosis in a sub-population of angiogenic cells. As used herein, the phrase "angiogenic cells" refers to any cells, which participate or contribute to the process of angiogenesis. Thus, angiogenic cells include but are not limited to, endothelial cells, smooth muscle cells.

In order to direct specific expression of an apoptotic agent in a subpopulation of angiogenic cells, the nucleic acid construct of the present invention includes a first polynucleotide region encoding a chimeric polypeptide including a ligand binding domain which can be, for example, a cell-surface receptor domain of a receptor tyrosine kinase, a receptor serine kinase, a receptor threonine kinase, a cell adhesion molecule or a phosphatase receptor fused to an effector domain of an apoptosis signaling molecule such as, for example, Fas, TNFR, and TRAIL.

Such a chimeric polypeptide can include any ligand binding domain fused to any apoptosis signaling domain as long as activation of the ligand binding domain, i.e., via ligand binding, triggers apoptosis signaling via the effector domain of the apoptosis signaling molecule.

Selection of the ligand binding domain and the apoptosis signaling domain fused thereto is affected according to the type of angiogenic cell targeted for apoptosis. For example, when targeting specific subset of endothelial cells (e.g., proliferating endothelial cells, or endothelial cells exhibiting a tumorous phenotype), the chimeric polypeptide includes a ligand binding domain capable of binding a ligand naturally present in the environment of such endothelial cells and preferably not present in endothelial cells of other non-targeted tissues (e.g., TNF, VEGF). Such a ligand can be secreted by endothelial cells (autocrine), secreted by neighboring tumor cells (paracrine) or specifically targeted to these endothelial cells.

Examples of suitable chimeric polypeptides are provided in Examples 2 of the Examples section which follows. Preferably, the chimeric polypeptide is the Fas-c chimera which is described in detail in Examples 2-4 of the Examples section which follows.

The use of such a chimeric polypeptide is particularly advantageous, since, as shown in the Examples section hereinunder, it enables efficient and robust activation of apoptosis in a specific subset of angiogenic cells while

avoiding activation in other subset of cells, which are not targeted for apoptosis.

To further enhance cell specificity of apoptosis, the nucleic acid construct of the present invention further includes a second polynucleotide region, which encodes a cis acting regulatory element (e.g., promoter and/or enhancer) capable of directing expression of the chimeric polypeptide in the sub-population of angiogenic cells.

Examples of suitable promoters/enhancers which can be utilized by the nucleic acid construct of the present invention include the endothelial-specific promoters: preproendothelin-1, PPE-1 promoter (Harats D, J Clin Invest. 1995 Mar;95(3):1335-44)., the PPE-1-3x promoter [PCT/IL01/01059; Varda-Bloom N, Gene Ther 2001 Jun;8(11):819-27], the TIE-1 (S79347, S79346) and the TIE-2 (U53603) promoters [Sato TN, Proc Natl Acad Sci U S A 1993 Oct 15;90(20):9355-8], the Endoglin promoter [Y11653; Rius C, Blood 1998 Dec 15;92(12):4677-90], the von Willerbrand factor [AF152417; Collins CJ Proc Natl Acad Sci U S A 1987 Jul;84(13):4393-7], the KDR/flk-1 promoter [X89777, X89776; Ronicke V, Circ Res 1996 Aug;79(2):277-85], The FLT-1 promoter [D64016 AJ224863; Morishita K, : J Biol Chem 1995 Nov 17;270(46):27948-53], the Egr-1 promoter [AJ245926; Sukhatme VP, Oncogene Res 1987 Sep-Oct;1(4):343-55], the E-selectin promoter [Y12462; Collins T J Biol Chem 1991 Feb 5;266(4):2466-73], The endothelial adhesion molecules promoters: ICAM-1 [X84737; Horley KJ EMBO J 1989 Oct;8(10):2889-96], VCAM-1 [M92431; Iademarco MF, J Biol Chem 1992 Aug 15;267(23):16323-9], PECAM-1 [AJ313330 X96849; CD31, Newman PJ, Science 1990 Mar 9;247(4947):1219-22], the vascular smooth-muscle-specific elements: CArG box X53154 and aortic carboxypeptidase-like protein (ACLP) promoter [AF332596; Layne MD, Circ Res. 2002; 90: 728-736] and Aortic Preferentially Expressed Gene-1 [Yen-Hsu Chen J. Biol. Chem., Vol. 276, Issue 50, 47658-47663, December 14, 2001].

Preferably, the promoter utilized by the construct of the present invention is functional in proliferating angiogenic cells, or angiogenic cells of a particular phenotype (e.g., tumorous). A promoter highly suitable for use with the present invention is the PPE-1-3x promoter which is further described in the Examples section which follows. A vector construct most suitable for use with the present invention is the widely used adenoviral vector and its derivatives.

The nucleic acid construct of the present invention can further include additional polynucleotide sequences such as for example, sequences encoding selection markers or reporter polypeptides, sequences encoding origin of replication in bacteria, sequences that allow for translation of several proteins from a single mRNA (IRES), sequences for genomic integration of the promoter-chimeric polypeptide encoding region and/or sequences generally included in mammalian expression vector such as pcDNA3, pcDNA3.1(+/-), pZeoSV2(+/-), pSecTag2, pDisplay, pEF/myc/cyto, pCMV/myc/cyto, pCR3.1, which are available from Invitrogen, pCI which is available from Promega, pBK-RSV and pBK-CMV which are available from Stratagene, pTRES which is available from Clontech, and their derivatives.

Preferably, the nucleic acid construct of the present invention is administered to the subject via, for example, systemic administration routes or via oral, rectal, transmucosal (especially transnasal), intestinal or parenteral administration routes. Systemic administration includes intramuscular, subcutaneous and intramedullary injections as well as intrathecal, direct intraventricular, intravenous, intraperitoneal, intranasal, intraocular injections or intra-tumoral.

Preferably, the subject is a mammal, more preferably, a human being, most preferably, a human being suffering from diseases characterized by excessive or abnormal neovascularization such as that characterizing tumor growth, proliferating diabetic retinopathy, arthritis and the like.

The nucleic acid constructs of the present invention can be administered to the subject *per se* or as part (active ingredient) of a pharmaceutical composition.

The prior art teaches of a number of delivery strategies which can be used to efficiently deliver naked or carrier provided polynucleotides into a wide variety of cell types (see, for example, Luft (1998) J Mol Med 76(2): 75-6; Kronenwett et al. (1998) Blood 91(3): 852-62; Rajur et al. (1997) Bioconjug Chem 8(6): 935-40; Lavigne et al. (1997) Biochem Biophys Res Commun 237(3): 566-71 and Aoki et al. (1997) Biochem Biophys Res Commun 231(3): 540-5).

As used herein a "pharmaceutical composition" refers to a preparation of one or more of the active ingredients or agents described herein with other chemical components such as physiologically suitable carriers and excipients. The purpose of a pharmaceutical composition is to facilitate administration of a compound to an organism.

Hereinafter, the phrases "physiologically acceptable carrier" and "pharmaceutically acceptable carrier" which may be interchangeably used refer to a carrier or a diluent that does not cause significant irritation to an organism and does not abrogate the biological activity and properties of the administered nucleic acid construct. An adjuvant is included under these phrases.

Herein the term "excipient" refers to an inert substance added to a pharmaceutical composition to further facilitate administration of an active ingredient. Examples, without limitation, of excipients include calcium carbonate, calcium phosphate, various sugars and types of starch, cellulose derivatives, gelatin, vegetable oils and polyethylene glycols.

Techniques for formulation and administration of drugs may be found in "Remington's Pharmaceutical Sciences," Mack Publishing Co., Easton, PA, latest edition, which is incorporated herein by reference.

Suitable routes of administration may, for example, include oral, rectal, transmucosal, especially transnasal, intestinal or parenteral delivery, including intramuscular, subcutaneous and intramedullary injections as well as intrathecal, direct intraventricular, intravenous, intraperitoneal, intranasal, or intraocular injections.

Alternately, one may administer the pharmaceutical composition in a local rather than systemic manner, for example, via injection of the pharmaceutical composition directly into a tissue region of a patient. In the context of the present invention, administration directly into tumor tissue is a relevant example of local administration.

Pharmaceutical compositions of the present invention may be manufactured by processes well known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilizing processes.

Pharmaceutical compositions for use in accordance with the present invention thus may be formulated in conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries, which facilitate processing of the active ingredients into preparations which can be used pharmaceutically. Proper formulation is dependent upon the route of administration chosen.

For injection, the active ingredient of the pharmaceutical composition may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hank's solution, Ringer's solution, or physiological salt buffer. For transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

For oral administration, the pharmaceutical composition can be formulated readily by combining the active compounds with pharmaceutically acceptable carriers well known in the art. Such carriers enable the pharmaceutical composition to be formulated as tablets, pills,

dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for oral ingestion by a patient. Pharmacological preparations for oral use can be made using a solid excipient, optionally grinding the resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries if
5 desired, to obtain tablets or dragee cores. Suitable excipients are, in particular, fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; cellulose preparations such as, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethyl-cellulose, sodium carbomethylcellulose; and/or
10 physiologically acceptable polymers such as polyvinylpyrrolidone (PVP). If desired, disintegrating agents may be added, such as cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate.

Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions may be used which may optionally contain gum
15 arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, titanium dioxide, lacquer solutions and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for identification or to characterize different combinations of active compound doses.

20 Pharmaceutical compositions which can be used orally, include push-fit capsules made of gelatin as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules may contain the active ingredients in admixture with filler such as lactose, binders such as starches, lubricants such as talc or magnesium stearate and,
25 optionally, stabilizers. In soft capsules, the active ingredients may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers may be added. All formulations for oral administration should be in dosages suitable for the chosen route of administration.

For buccal administration, the compositions may take the form of tablets or lozenges formulated in conventional manner.

For administration by nasal inhalation, the active ingredients for use according to the present invention are conveniently delivered in the form of
5 an aerosol spray presentation from a pressurized pack or a nebulizer with the use of a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichloro-tetrafluoroethane or carbon dioxide. In the case of a pressurized aerosol, the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of,
10 e.g., gelatin for use in a dispenser may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

The pharmaceutical composition described herein may be formulated for parenteral administration, e.g., by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, e.g., in
15 ampoules or in multidose containers with optionally, an added preservative. The compositions may be suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents.

Pharmaceutical compositions for parenteral administration include
20 aqueous solutions of the active preparation in water-soluble form. Additionally, suspensions of the active ingredients may be prepared as appropriate oily or water based injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acids esters such as ethyl oleate, triglycerides or liposomes. Aqueous
25 injection suspensions may contain substances, which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol or dextran. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the active ingredients to allow for the preparation of highly concentrated solutions.

Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, e.g., sterile, pyrogen-free water based solution, before use.

The pharmaceutical composition of the present invention may also be
5 formulated in rectal compositions such as suppositories or retention enemas, using, e.g., conventional suppository bases such as cocoa butter or other glycerides.

Pharmaceutical compositions suitable for use in context of the present invention include compositions wherein the active ingredients are contained
10 in an amount effective to achieve the intended purpose. More specifically, a therapeutically effective amount means an amount of active ingredients (e.g. antisense oligonucleotide) effective to prevent, alleviate or ameliorate symptoms of a disorder (e.g., progressive loss of bone mass) or prolong the survival of the subject being treated.

15 Determination of a therapeutically effective amount is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein.

For any preparation used in the methods of the invention, the therapeutically effective amount or dose can be estimated initially from *in*
20 *vitro* and cell culture assays. For example, a dose can be formulated in an animal model, such as the murine Src deficient model of osteopetrosis (Boyce et al. (1992) J. Clin. Invest. 90, 1622-1627; Lowe et al. (1993) Proc. Natl. Acad. Sci. USA 90, 4485-4489; Soriano et al. (1991) Cell 64, 693-702), to achieve a desired concentration or titer. Such information can be
25 used to more accurately determine useful doses in humans.

Toxicity and therapeutic efficacy of the active ingredients described herein can be determined by standard pharmaceutical procedures *in vitro*, in cell cultures or experimental animals. The data obtained from these *in vitro* and cell culture assays and animal studies can be used in formulating a range
30 of dosage for use in human. The dosage may vary depending upon the

dosage form employed and the route of administration utilized. The exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition. (See e.g., Fingl, *et al.*, 1975, in "The Pharmacological Basis of Therapeutics", Ch. 1 p.1).

- 5 Dosage amount and interval may be adjusted individually to levels of the active ingredient are sufficient to retard tumor progression (minimal effective concentration, MEC). The MEC will vary for each preparation, but can be estimated from *in vitro* data. Dosages necessary to achieve the MEC will depend on individual characteristics and route of administration.
- 10 Detection assays can be used to determine plasma concentrations.

Depending on the severity and responsiveness of the condition to be treated, dosing can be of a single or a plurality of administrations, with course of treatment lasting from several days to several weeks or diminution of the disease state is achieved.

- 15 The amount of a composition to be administered will, of course, be dependent on the subject being treated, the severity of the affliction, the manner of administration, the judgment of the prescribing physician, etc.

- Compositions of the present invention may, if desired, be presented in a pack or dispenser device, such as an FDA approved kit, which may contain
- 20 one or more unit dosage forms containing the active ingredient. The pack may, for example, comprise metal or plastic foil, such as a blister pack. The pack or dispenser device may be accompanied by instructions for administration. The pack or dispenser may also be accommodated by a notice associated with the container in a form prescribed by a governmental
- 25 agency regulating the manufacture, use or sale of pharmaceuticals, which notice is reflective of approval by the agency of the form of the compositions or human or veterinary administration. Such notice, for example, may be of labeling approved by the U.S. Food and Drug Administration for prescription drugs or of an approved product insert. Compositions comprising a
- 30 preparation of the invention formulated in a compatible pharmaceutical

carrier may also be prepared, placed in an appropriate container, and labeled for treatment of an indicated condition, as if further detailed above.

The pharmaceutical compositions of the present invention may further include any additional ingredients which may improve the uptake of the nucleic acid construct by the cells, expression of the chimeric polypeptide encoded by the nucleic acid construct in the cells, or the activity of the expressed chimeric polypeptide.

For example, the uptake of adenoviral vectors into EC cells can be enhanced by treating the vectors with engineered antibodies or small peptides. Such "adenobody" treatment, was shown effective in directing adenovirus constructs to EGF receptors on cells (Watkins et al 1997, Gene Therapy 4:1004-1012). In addition, Nicklin et al have shown that A small peptide, isolated via phage display, increased specificity and efficiency of vectors in endothelial cells and decreased the expression in liver cells in culture (Nicklin et al 2000, Circulation 102:231-237). In a recent study, an FGF retargeted adenoviral vector reduced the toxicity of tk in mice (Printz et al 2000, Human Gene Therapy 11:191-204).

It will be appreciated that although targeting of cells exposed to the ligand is preferred, the present invention also envisages expression of the nucleic acid construct of the present invention in cells which are not exposed to, or naturally affected by the ligand. In such cases, the method of the present invention includes the step of administering such a ligand to the cells transformed. Such administration can be effected by using any of the above described administration methods. Preferably, the ligand is administered in a cell targeted manner, using for example antibody conjugated targeting, such that activation of apoptosis signaling is highly specific. This approach of apoptosis activation is described in more detail in the Examples section which follows.

Thus, the present invention provides nucleic acid constructs, pharmaceutical compositions including such constructs and methods of

utilizing such constructs for down-regulating angiogenesis in tissue regions characterized by excessive or abnormal angiogenesis.

Since the present invention enables targeted expression in specific cell subsets, it can also be modified and used in for treating various tumors.

5 Thus, according to another aspect of the present invention there is provided a method of treating tumors.

The method according to this aspect of the present invention is effected by expressing in tumor cells the chimeric polypeptide described above.

10 Thus according to this aspect of the present invention, expression of the polypeptide chimera is directed by a tumor specific element, such as, but not limited to, the gastrin-releasing peptide (GRP) promoter [AF293321S3; Morimoto E Anticancer Res 2001 Jan-Feb;21(1A):329-31], the hTERT promoter [AH007699; Gu J , Gene Ther 2002 Jan;9(1):30-7], the Hexokinase
15 type II promoter [AF148512; Katabi MM, Hum Gene Ther. 1999 Jan 20;10(2):155-64.], or the L-plastin promoter [L05490, AH002870, MMU82611; Peng XY, Cancer Res. 2001 Jun 1;61(11):4405-13].

Expression of the polypeptide chimera (e.g., Fas-c) in tumor cells activates apoptosis in these cells and thus leads to cell death, which in turn
20 causes tumor growth slowdown or arrest, and possibly tumor shrinkage.

Additional objects, advantages, and novel features of the present invention will become apparent to one ordinarily skilled in the art upon examination of the following examples, which are not intended to be limiting.
25 Additionally, each of the various embodiments and aspects of the present invention as delineated hereinabove and as claimed in the claims section below finds experimental support in the following examples.

EXAMPLES

Reference is now made to the following examples, which together with the above descriptions, illustrate the invention in a non limiting fashion.

Generally, the nomenclature used herein and the laboratory procedures in recombinant DNA technology described below are those well known and commonly employed in the art. Standard techniques are used for cloning, DNA and RNA isolation, amplification and purification. Generally enzymatic reactions involving DNA ligase, DNA polymerase, restriction endonucleases and the like are performed according to the manufacturers' specifications. These techniques and various other techniques are generally performed according to Sambrook et al., Molecular Cloning--A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y. (1989). The manual is hereinafter referred to as "Sambrook". Other general references are provided throughout this document. The procedures therein are believed to be well known in the art and are provided for the convenience of the reader. All the information contained therein is incorporated herein by reference.

EXAMPLE 1

***In-vitro assay for pro-apoptotic gene activity in endothelial cells
(BAEC) and 293 cells***

In cancer treatment, anti-angiogenic therapy targets the evolving vasculature which nourishes the growing tumor [Folkman J. N Engl J Med (1995) 333(26):1757-63]. As the research of apoptosis, or programmed cell death, has progressed, numerous genes that encode selective and efficient cell death regulators have been identified [Strasser et al. Annu Rev Biochem (2000) 69:217-45.].

The present study screened several pro-apoptotic genes in order to identify an agent most suitable for anti-angiogenic therapy. Several pro-apoptotic genes including MORT1 (FADD - Fas associated death domain protein, GenBank Accession number NM_003824), RIP (receptor-interacting-

protein, GenBank Accession number U25995), CASH (c-FLIP, GenBank
Accession number AF010127), MACH (caspase 8 GenBank Accession
number X98172), CPP32 (caspase 3, GenBank Accession number U13737),
caspase 9 (U60521) and Fas-chimera (Fas-c), a previously described fusion of
5 two "death receptors", constructed from the extracellular region of TNFR1
and the trans-membrane and intracellular regions of Fas [Boldin MP et al. J
Biol Chem (1995) 270(14):7795-8, see Figure 1a) were PCR amplified and
cloned into the pcDNA3 (Invitrogen, Inc.) mammalian expression vector
using well known prior art cloning techniques.

10 These pro-apoptotic gene constructs were co-expressed along with
pGFP in BAEC (Bovine Aortic Endothelial Cells) and 293 cells, which were
used as non-endothelial control cells. 24 hours post transfection, cells were
analyzed using fluorescent microscopy. Apoptotic cells were identified based
on typical morphology, (i.e., small and round shape) using fluorescence
15 microscopy (Figures 2a-b). Further assessment of the apoptotic phenotype
was effected using electron microscopy (Figures 3a-f). Quantification of the
apoptotic effect showed that MORT1, TNFR1 and Fas-chimera induced the
highest apoptotic activity in BAEC and 293 cells (Figure 4a-b). Caspase 3
and 9 were less potent in this respect, probably because they were in an
20 inactive zymogen form. Based on these results, the Fas-chimera (Fas-c) gene
was selected for the generation of an adenoviral-vector to be used in anti-
angiogenic therapy.

EXAMPLE 2

25 *Production of recombinant adenoviruses encoding Fas-chimera under the control of the modified PPE-1 promoter*

A cDNA encoding a full length Fas-chimera was subcloned into the
plasmid pPACPPE1-3x containing the modified pre-proendothelin1 promoter
(see Figure 1b). Recombinant adenoviruses were produced by co-transfection

of this plasmid with pJM17 plasmid into human embryonic kidney 293 cells. Successful viral cloning was verified via PCR amplification (Figure 5a).

In order to determine the expression of Fas-c in the target cells, endothelial BAEC cells were transduced with the indicated titer of Ad-PPE-Fas-c. 72h post transduction cells were lysed and cellular proteins resolved using a non-reducing SDS-PAGE gel. Western blot analysis was performed using anti-TNFR1 antibody (Sc-7895, Santa-Cruz Biotech). As demonstrated in Figure 5b, a prominent band migrating at 45kD was clearly evident and its expression was dose-dependent, suggesting correct folding and expression of the chimeric protein. In contrast, no corresponding bands were evident in non-transduced endothelial cells or in cells transduced with control empty viral vector. Thus, these results confirmed that the adenoviral-mediated gene transfer of Fas-c results in transgene expression in the target cells.

EXAMPLE 3

Ad-PPE-Fas-c expression induces apoptosis in endothelial cells

The ability of Ad-PPE-Fas chimera to induce apoptosis of endothelial cells was determined. As shown in Figures 6a-b, pre-proendothelin directed, adenovirus-mediated transduction of endothelial cells resulted in an evident and massive cell death; HUVEC and BAEC infected with Ad-PPE-Fas-c (10^3 MOI) had morphological features of adherent cells undergoing apoptosis including membrane blebbing, rounding and shrinking and detachment from the culture dish. In contrast, cells infected with control viruses at the same MOI, maintained normal appearance and growth rate. Cells transduced with 100 MOI presented only a minimal degree of cell death (data not shown).

Further assessment of the cytotoxic properties of Ad-PPE-Fas-c was effected by expressing this virus in cells expressing the reporter gene GFP under the control of the PPE-1 promoter. As is evident from Figures 6c-d, most of the transduced cells acquired a typical apoptotic appearance 72 hours

following transduction, whereas cells co-transduced with control virus and Ad-PPE-GFP appeared normal.

The cytotoxic effect of Fas-c was quantified using crystal violet staining. As shown in Figure 7, infection of BAEC and HUVEC with Ad-PPE-Fas-c resulted in mortality rates of 57 % and 65 %, respectively, while the control virus did not affect cell viability.

The endothelial cell specificity of the pro-apoptotic vector Ad-PPE-Fas- was demonstrated by infecting NSF (normal skin fibroblasts) with this vector. These cells, which express low levels of PPE-1 [Varda-Bloom, N. et al. Gene Ther 8, 819-27. (2001)] were not affected by infection with Ad-PPE-Fas-c. In contrast, the recombinant vector Ad-CMV-Fas-c induced apoptotic in these cells.

EXAMPLE 4

Co-administration of Ad-PPE-Fas-c receptor and TNF α ligand augments the pro-apoptotic effect in a selective manner

The ability of TNF α to augment the apoptotic effect in Fas-c expressing cells was investigated. Human TNF α was added to an endothelial cell culture 48h-post virus infection with Ad-PPE-Fas-c (MOI of 100). Cell viability was assayed 24 h later. As shown in Figure 8, TNF α (10ng/ml) induced a 73% decrease in viability of Ad-PPE-Fas-c infected cells, whereas no significant mortality was effected by TNF α alone or in cells infected with control virus (Ad-Luc).

To substantiate the effect of TNF α , cell specificity was addressed. NSF (normal skin fibroblasts), DA3 (mouse mammary adenocarcinoma), D122 (Lewis lung carcinoma) and B16 melanoma cells were infected with Ad-PPE-Fas-c or a control virus. 48 hours later, culture was supplemented with TNF α and cell morphology was assessed following staining with crystal violet. As shown in Figures 9a-e, non-endothelial cells infected with Ad-PPE-Fas-c displayed normal appearance and were not affected by TNF. On

the other hand, adenoviral mediated infection of BAEC with Fas-c resulted in marked decrease in cell viability when TNF was added. The non-selective apoptotic activity of Fas-c driven by CMV promoter is demonstrated in Figure 10a which illustrates the TNF-dependent apoptotic effect of Ad-CMV-Fas-c on endothelial cells. Viability of BAEC cells infected with the indicated MOI of Ad-CMV-Fas-chimera was determined following incubation with TNF.

Notably, the non-endothelial-specific vector Ad-CMV-Fas-c caused TNF α -dependent apoptosis of both endothelial and non-endothelial cells (Figures 10b-d).

EXAMPLE 5

Ad-PPE1-Fas-c induces in-vivo growth retardation of B16 melanoma in mice

The B16 melanoma mouse model was used in order to test the anti-tumoral effect of Fas-c expressed from the PPE1-3x promoter. B16 melanoma cells (8×10^5) were injected subcutaneously to the flank region of 40 C57bl/6 mice. When the tumor was palpable ($\approx 5 \times 5$ mm), the mice were randomized into 4 groups as follows: (i) control- saline injection; (ii) control virus (Adeno virus containing luciferase controlled by PPE promoter); (iii) Ad-PPE1-3x-Fas-c- virus containing the Fas -TNF receptor chimeric gene controlled by the preproendothelin (PPE) promoter; and (iv) Ad-CMV- Fas-c- virus containing the Fas -TNF receptor chimeric gene controlled by the non-endothelial specific, CMV promoter.

Tumor size (length and width) was measured using a hand caliper. As shown in Figure 11a, tumor size was lower for mice treated with Ad-PPE1-3x-Fas-c or Ad-CMV-Fas-c as compared to control mice. Tumor weights at the end of the treatment period was also lower in the Ad-PPE1-3x-Fas-c treated mice (Figure 11b). Mice injected with Ad-PPE1-3x-Fas-c showed marked necrosis of their tumor (Figure 11c).

Although the invention has been described in conjunction with specific embodiments thereof, it is evident that many alternatives, modifications and variations will be apparent to those skilled in the art. Accordingly, it is intended to embrace all such alternatives, modifications and variations that
5 fall within the spirit and broad scope of the appended claims. All publications, patents, patent applications and sequences identified by their accession numbers mentioned in this specification are herein incorporated in their entirety by reference into the specification, to the same extent as if each individual publication, patent, patent application or sequence identified by
10 their accession number was specifically and individually indicated to be incorporated herein by reference. In addition, citation or identification of any reference in this application shall not be construed as an admission that such reference is available as prior art to the present invention.

WHAT IS CLAIMED IS:

1. A nucleic acid construct comprising:
 - (a) a first polynucleotide region encoding a chimeric polypeptide including a ligand binding domain fused to an effector domain of an apoptosis signaling molecule; and
 - (b) a second polynucleotide region encoding a cis acting regulatory element being for directing expression of said chimeric polypeptide in a specific tissue or cell;

wherein said ligand binding domain is selected such that it is capable of binding a ligand present in said specific tissue or cell, whereas binding of said ligand to said ligand binding domain activates said effector domain of said apoptosis signaling molecule.

2. The nucleic acid construct of claim 1, wherein said cis acting regulatory element is an endothelial cell-specific or periendothelial cell-specific promoter selected from the group consisting of the PPE-1 promoter, the PPE-1-3x promoter, the TIE-1 promoter, the TIE-2 promoter, the Endoglin promoter, the von Willerband promoter, the KDR/flk-1 promoter, The FLT-1 promoter, the Egr-1 promoter, the ICAM-1 promoter, the VCAM-1 promoter, the PECAM-1 promoter and the aortic carboxypeptidase-like protein (ACLP) promoter.

3. The nucleic acid construct of claim 1, wherein said ligand binding domain is a ligand-binding domain of a cell-surface receptor.

4. The nucleic acid construct of claim 3, wherein said cell-surface receptor is selected from the group consisting of a receptor tyrosine kinase, a receptor serine kinase, a receptor threonine kinase, a cell adhesion molecule and a phosphatase receptor.

5. The nucleic acid construct of claim 1, wherein said apoptosis signaling molecule is selected from the group consisting of Fas, TNFR, and TRAIL.

6. A mammalian cell transformed with the nucleic acid construct of claim 1.

7. A method of down-regulating angiogenesis in a tissue of a subject, the method comprising administering to the subject a nucleic acid construct designed and configured for generating apoptosis in a sub-population of angiogenic cells, said nucleic acid construct including:

- (a) a first polynucleotide region encoding a chimeric polypeptide including a ligand binding domain fused to an effector domain of an apoptosis signaling molecule; and
- (b) a second polynucleotide region encoding a cis acting regulatory element being for directing expression of said chimeric polypeptide in said sub-population of angiogenic cells;

wherein said ligand binding domain is selected such that it is capable of binding a ligand present in, or provided to, said sub-population of angiogenic cells, whereas binding of said ligand to said ligand binding domain activates said effector domain of said apoptosis signaling molecule, thereby down-regulating angiogenesis in the tissue.

8. The method of claim 7, further comprising administering said ligand to the subject in a manner suitable for providing said ligand to said sub-population of angiogenic cells.

9. The method of claim 7, wherein said administering said ligand to the subject is effected by a method selected from the group consisting of:

- (i) systemic in-vivo administration;

- (ii) ex-vivo administration to cells removed from a body of the subject, said cells subsequently reintroduced into said body of the subject; and
- (iii) local in-vivo administration.

10. The method of claim 7, wherein said cis acting regulatory element is an endothelial cell-specific or periendothelial cell-specific promoter selected from the group consisting of the PPE-1 promoter, the PPE-1-3x promoter, the TIE-1 promoter, the TIE-2 promoter, the Endoglin promoter, the von Willerband promoter, the KDR/flk-1 promoter, The FLT-1 promoter, the Egr-1 promoter, the ICAM-1 promoter, the VCAM-1 promoter, the PECAM-1 promoter and the aortic carboxypeptidase-like protein (ACLP) promoter.

11. The method of claim 7, wherein said ligand binding domain is a ligand-binding domain of a cell-surface receptor.

12. The method of claim 11, wherein said cell-surface receptor is selected from the group consisting of a receptor tyrosine kinase, a receptor serine kinase, a receptor threonine kinase, a cell adhesion molecule and a phosphatase receptor.

13. The method of claim 7, wherein said apoptosis signaling molecule is selected from the group consisting of Fas, TNFR and TRAIL.

14. A method of down-regulating angiogenesis in a tissue of a subject, the method comprising:

- (a) administering to the subject a nucleic acid construct designed and configured for generating apoptosis in a sub-population of angiogenic cells, said nucleic acid construct including:

- (i) a first polynucleotide region encoding a chimeric polypeptide including a ligand binding domain fused to an effector domain of an apoptosis signaling molecule, wherein said effector domain is selected such that it is activated following binding of a ligand to said ligand binding domain; and
 - (ii) a second polynucleotide region encoding a cis acting regulatory element being for directing expression of said chimeric polypeptide in said sub-population of angiogenic cells; and
- (b) administering to the subject said ligand, thereby down-regulating angiogenesis in the tissue.

15. The method of claim 14, wherein said administering said ligand to the subject is effected by a method selected from the group consisting of:

- (i) systemic in-vivo administration;
- (ii) ex-vivo administration to cells removed from a body of the subject, said cells subsequently reintroduced into said body of the subject; and
- (iii) local in-vivo administration.

16. The method of claim 14, wherein said cis acting regulatory element is an endothelial cell-specific or periendothelial cell-specific promoter selected from the group consisting of PPE-1 promoter, the PPE-1-3x promoter, the TIE-1 promoter, the TIE-2 promoter, the Endoglin promoter, the von-Willebrand promoter, the KDR/flk-1 promoter, The FLT-1 promoter, the Egr-1 promoter, the ICAM-1 promoter, the VCAM-1 promoter, the PECAM-1 promoter and the aortic carboxypeptidase-like protein (ACLP) promoter.

17. The method of claim 14, wherein said ligand binding domain is a ligand-binding domain of a cell-surface receptor.

18. The method of claim 17, wherein said cell-surface receptor is selected from the group consisting of a receptor tyrosine kinase, a receptor serine kinase, a receptor threonine kinase, a cell adhesion molecule and a phosphatase receptor.

19. The method of claim 14, wherein said apoptosis signaling molecule is selected from the group consisting of Fas, TNFR and TRAIL.

20. A pharmaceutical composition for down regulating angiogenesis in a tissue of a subject, the pharmaceutical composition comprising as an active ingredient a nucleic acid construct designed and configured for generating apoptosis in a subpopulation of angiogenic cells and a pharmaceutical acceptable carrier, said nucleic acid construct including:

- (a) a first polynucleotide region encoding a chimeric polypeptide including a ligand binding domain fused to an effector domain of an apoptosis signaling molecule; and
- (b) a second polynucleotide region encoding a cis acting regulatory element being for directing expression of said chimeric polypeptide in said subpopulation of angiogenic cells;

wherein said ligand binding domain is selected such that it is capable of binding a ligand present in said specific tissue or cell, whereas binding of said ligand to said ligand binding domain activates said effector domain of said apoptosis signaling molecule.

21. The pharmaceutical composition of claim 20, wherein said cis acting regulatory element is an endothelial cell-specific or periendothelial cell-specific promoter selected from the group consisting of the PPE-1

promoter, the PPE-1-3x promoter, the TIE-1 promoter, the TIE-2 promoter, the Endoglin promoter, the von Willerband promoter, the KDR/flk-1 promoter, The FLT-1 promoter, the Egr-1 promoter, the ICAM-1 promoter, the VCAM-1 promoter, the PECAM-1 promoter and the aortic carboxypeptidase-like protein (ACLP) promoter.

22. The pharmaceutical composition of claim 20, wherein said ligand binding domain is a ligand-binding domain of a cell-surface receptor.

23. The pharmaceutical composition of claim 22, wherein said cell-surface receptor is selected from the group consisting of a receptor tyrosine kinase, a receptor serine kinase, a receptor threonine kinase, a cell adhesion molecule and a phosphatase receptor.

24. The pharmaceutical composition of claim 20, wherein said apoptosis signaling molecule is selected from the group consisting of Fas, TNFR and TRAIL.

25. A method of treating a disease or condition associated with excessive neo-vascularization, the method comprising administering a therapeutically effective amount of a nucleic acid construct designed and configured for generating apoptosis in a sub-population of angiogenic cells, said nucleic acid construct including:

- (i) a first polynucleotide region encoding a chimeric polypeptide including a ligand binding domain fused to an effector domain of an apoptosis signaling molecule; and
- (ii) a second polynucleotide region encoding a cis acting regulatory element being for directing expression of said chimeric polypeptide in said sub-population of angiogenic cells;

wherein said ligand binding domain is selected such that it is capable of binding a ligand present in, or provided to, said sub-population of angiogenic cells, whereas binding of said ligand to said ligand binding domain activates said effector domain of said apoptosis signaling molecule, thereby down-regulating angiogenesis in the tissue and treating the disease or condition associated with excessive neo-vascularization.

26. The method of claim 25, further comprising administering said ligand to the subject in a manner suitable for providing said ligand to said sub-population of angiogenic cells.

27. The method of claim 26, wherein said administering said ligand to the subject is effected by a method selected from the group consisting of:

- (i) systemic in-vivo administration;
- (ii) ex-vivo administration to cells removed from a body of the subject, said cells subsequently reintroduced into said body of the subject; and
- (iii) local in-vivo administration.

28. The method of claim 25, wherein said cis acting regulatory element is an endothelial cell-specific or periendothelial cell-specific promoter selected from the group consisting of the PPE-1 promoter, the PPE-1-3x promoter, the TIE-1 promoter, the TIE-2 promoter, the Endoglin promoter, the von Willerband promoter, the KDR/flk-1 promoter, The FLT-1 promoter, the Egr-1 promoter, the ICAM-1 promoter, the VCAM-1 promoter, the PECAM-1 promoter and the aortic carboxypeptidase-like protein (ACLP) promoter.

29. The method of claim 25, wherein said ligand binding domain is a ligand-binding domain of a cell-surface receptor.

30. The method of claim 29, wherein said cell-surface receptor is selected from the group consisting of a receptor tyrosine kinase, a receptor serine kinase, a receptor threonine kinase, a cell adhesion molecule and a phosphatase receptor.

31. The method of claim 25, wherein said apoptosis signaling molecule is selected from the group consisting of Fas, TNFR and TRAIL.

32. A method of treating a tumor in a subject, the method comprising administering a therapeutically effective amount of a nucleic acid construct designed and configured for generating apoptosis in cells of the tumor, said nucleic acid construct including:

- (i) a first polynucleotide region encoding a chimeric polypeptide including a ligand binding domain fused to an effector domain of an apoptosis signaling molecule; and
- (ii) a second polynucleotide region encoding a cis acting regulatory element being for directing expression of said chimeric polypeptide in said cells of the tumor;

wherein said ligand binding domain is selected such that it is capable of binding a ligand present in, or provided to, said cells of the tumor, whereas binding of said ligand to said ligand binding domain activates said effector domain of said apoptosis signaling molecule to thereby direct apoptosis in said cells of the tumor.

33. The method of claim 32, further comprising administering said ligand to the subject in a manner suitable for providing said ligand to said cells of the tumor.

34. The method of claim 32, wherein said cis acting regulatory element is selected from the group consisting of the gastrin-releasing peptide

(GRP) promoter, the hTERT promoter, the Hexokinase type II promoter and the L-plastin promoter.

1/9




Fig. 1a




Fig. 1b



Fig. 2a



Fig. 2b

Fig. 3a



Fig. 3b




Fig. 3c





Fig. 3d




Fig. 3e



Fig. 3f

3/9




Fig. 4

4/9




Fig. 5a




Fig. 5b




Fig. 6a




Fig. 6b

5/9

Ad-PPE-Fas-c



Fig. 6c

Ad-PPE-Luc




Fig. 6d




Fig. 7

6/9






Fig. 8



BAEC

Fig. 9a



NSF

Fig. 9b

7/9




Fig. 9c




Fig. 9d




Fig. 9e

8/9




Fig. 10a




Fig. 10b

Fig. 10c

Fig. 10d

9/9




Fig. 11a




Fig. 11b




Fig. 11c

INTERNATIONAL SEARCH REPORT

International application No.

PCT/IL02/00339

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) :C07H 21/04; A61K 48/00

US CL :536/23.1; 514/44

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 536/23.1; 514/44

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
none

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

MEDLINE, WEST, BIOSIS, EMBASE

search terms: Harats D., enbrel, TNF r, TRAIL, apoptosis, angiogenesis, cis acting element, promoter,

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 6,180,355 B1 (ALEXANDER et al.) 30 January 2001, see entire document and column 9, lines 13-25.	1-34
Y	US 6,239,151 B1 (BROADHURST et al) 29 May 2001, abstract and see entire document.	1-34

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

16 SEPTEMBER 2002

Date of mailing of the international search report

02 DEC 2002

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

Michael A. Belyavsky

Telephone No. (703) 308-0196